arXiv:1510.07617v1 [math-ph] 26 Oct 2015

LOCALLY CONVEX QUASI C*-ALGEBRAS AND NONCOMMUTATIVE INTEGRATION

CAMILLO TRAPANI AND SALVATORE TRIOLO

Abstract. In this paper we continue the analysis undertaken in a series of previous papers on structures arising as completions of C*-algebras under topologies coarser that their norm and we focus our attention on the so-called *locally convex quasi C*-algebras*. We show, in particular, that any strongly *-semisimple locally convex quasi C*-algebra $(\mathfrak{X}, \mathfrak{A}_0)$, can be represented in a class of noncommutative local L^2-spaces.

1. Introduction

The completion $\mathfrak{X}$ of a C*-algebra $\mathfrak{A}_0$, with respect to a norm weaker than the C*-norm provides a mathematical framework where discussing certain quantum physical systems for which the usual algebraic approach made in terms of C*-algebras revealed to be insufficient.

First of all, $\mathfrak{X}$ is a Banach $\mathfrak{A}_0$-module and becomes a quasi *-algebra if $\mathfrak{X}$ carries an involution which extends the involution $*$ of $\mathfrak{A}_0$. This structure has been called a *proper CQ*-algebra* in a series of papers [4]-[10], [22]-[21] to that we refer for a detailed analysis. On the other hand, if $\mathfrak{X}$ is endowed with an isometric involution different from that of $\mathfrak{A}_0$, then the structure becomes more involved.

CQ*-algebras are examples of more general structures called *locally convex quasi C*-algebras* [3]. They are obtained by completing a C*-algebra with respect to a new locally convex topology τ on $\mathfrak{A}_0$ compatible with the corresponding $\|\cdot\|$-topology. Under certain conditions on τ a quasi $*$-subalgebra $\mathfrak{A}$ of the completion $\widetilde{\mathfrak{A}_0}[\tau]$ is a locally convex quasi *-algebra which is named locally convex quasi C*-algebra.

In [9] quasi *-algebras of measurable and/or integrable operators (in the sense of Segal [19], [27] and Nelson [17]) were examined in detail and it was proved that any *-semisimple CQ*-algebra can be realized as a CQ*-algebra of measurable operators, with the help of a particular class of positive bounded sesquilinear forms on $\mathfrak{X}$.

In this paper, after a short overview of the main results obtained on this subject, we continue our study of locally convex quasi C*-algebras and we generalize the results obtained in [9] for proper CQ*- algebras to these structures.

The main question we pose now in the present paper is the following: given a *-semisimple locally convex quasi C*-algebras $(\mathfrak{X}, \mathfrak{A}_0)$ and the universal *-representation of $\mathfrak{A}_0$, defined via the Gelfand-Naimark theorem, can $\mathfrak{X}$ be realized as a locally convex quasi C*-algebra of operators of type L^2?

2000 *Mathematics Subject Classification.* Primary 46L08; Secondary 46L51, 47L60 .

Key words and phrases. Banach C*-modules, Noncommutative integration, Partial algebras of operators.

The paper is organized as follows. We begin with giving a short overview of noncommutative L^p-spaces (constructed starting from a von Neumann algebra $\mathfrak{M}$ and a normal, semifinite, faithful trace φ on $\mathfrak{M}$), considered as CQ*-algebras. We also introduce the noncommutative $L^p_{\rm loc}$-space constructed on a von Neumann algebra possessing a family of mutually orthogonal central projections whose sum is the identity operator. We show that $(L^p_{\rm loc}(\varphi), \mathfrak{M})$ is a locally convex quasi C*-algebra.

Finally we give some results on the structure of locally convex quasi C*-algebras: we prove that any locally convex quasi C*-algebra $(\mathfrak{X}, \mathfrak{A}_0)$ possessing a sufficient family of bounded positive tracial sesquilinear forms can be continuously embedded into a locally convex quasi C*-algebras of measurable operators of the type $(L^2_{\rm loc}(\varphi), \mathfrak{M})$.

1.1. **Definitions and results on noncommutative measure.** The following basic definitions and results on noncommutative measure theory and integration are needed in what follows. Let $\mathfrak{M}$ be a von Neumann algebra on a Hilbert space $\mathcal{H}$ and φ a normal faithful semifinite trace defined on $\mathfrak{M}_+$.

Put

$$\mathcal{J} = \{X \in \mathfrak{M} : \varphi(|X|) < \infty\}.$$

$\mathcal{J}$ is a *-ideal of $\mathfrak{M}$.

Let $P \in \mathrm{Proj}(\mathfrak{M})$, the lattice of projections of $\mathfrak{M}$. Two projections $P, Q \in \mathrm{Proj}(\mathfrak{M})$ are called equivalent, $P \sim Q$ if there is a $U \in \mathfrak{M}$ with $U^*U = P$ and $UU^* = Q$. We say that $P \prec Q$ in the case where P is equivalent to a subprojection of Q.

A projection P of a von Neumann algebra $\mathfrak{M}$ is said to be *finite* if $P \sim Q \leq P$ implies $P = Q$. A projection $P \in \mathfrak{M}$ is said to be *purely infinite* if there is no nonzero finite projection $Q \preceq P$ in $\mathfrak{M}$. A von Neumann algebra $\mathfrak{M}$ is said to be *finite* (respectively, *purely infinite*) if the identity operator $\mathbb{I}$ is finite (respectively, purely infinite).

We say that P is φ-finite if $P \in \mathcal{J}$. Any φ-finite projection is finite.

In what follows we will need the following result (see, [15, Vol. IV, Ex. 6.9.12]) that we state as a lemma.

Lemma 1.1. *Let $\mathfrak{M}$ be a von Neumann algebra on a Hilbert space $\mathcal{H}$ and φ a normal faithful semifinite trace defined on $\mathfrak{M}_+$. There is an orthogonal family $\{Q_j; j \in J\}$ of nonzero central projections in $\mathfrak{M}$ such that $\bigvee_{j\in J} Q_j = \mathbb{I}$ and each Q_j is the sum of an orthogonal family of mutually equivalent finite projections in $\mathfrak{M}$.*

A vector subspace $\mathcal{D}$ of $\mathcal{H}$ is said to be *strongly dense* (resp., *strongly φ-dense*) if

- $U'\mathcal{D} \subset \mathcal{D}$ for any unitary U' in $\mathfrak{M}'$;
- there exists a sequence $P_n \in \mathrm{Proj}(\mathfrak{M})$: $P_n\mathcal{H} \subset \mathcal{D}$, $P_n^\perp \downarrow 0$ and $P_n^\perp$ is a finite projection (resp., $\varphi(P_n^\perp) < \infty$).

Clearly, every strongly φ-dense domain is strongly dense.

Throughout this paper, when we say that an operator T is affiliated with a von Neumann algebra, written $T\,\eta\,\mathfrak{M}$, we always mean that T is closed, densely defined and $TU \supseteq UT$ for every unitary operator $U \in \mathfrak{M}'$.

An operator $T\,\eta\,\mathfrak{M}$ is called

- *measurable* (with respect to $\mathfrak{M}$) if its domain $D(T)$ is strongly dense;

- φ-*measurable* if its domain $D(T)$ is strongly φ-dense.

From the definition itself it follows that, if T is φ-measurable, then there exists $P \in \mathrm{Proj}(\mathfrak{M})$ such that TP is bounded and $\varphi(P^\perp) < \infty$.

We remind that any operator affiliated with a finite von Neumann algebra is measurable [19, Cor. 4.1] but it is not necessarily φ-measurable.

Remark 1.2. The following statements will be used later.

(i) Let $T \,\eta\, \mathfrak{M}$ and $Q \in \mathfrak{M}$. If $D(TQ) = \{\xi \in \mathcal{H} : Q\xi \in D(T)\}$ is dense in $\mathcal{H}$, then $TQ \,\eta\, \mathfrak{M}$.

(ii) If $Q \in \mathrm{Proj}(\mathfrak{M})$, then $Q\mathfrak{M}Q = \{QXQ \restriction_{Q\mathcal{H}}; X \in \mathfrak{M}\}$ is a von Neumann algebra on the Hilbert space $Q\mathcal{H}$; moreover $(Q\mathfrak{M}Q)' = Q\mathfrak{M}'Q$. If $T \,\eta\, \mathfrak{M}$ and $Q \in \mathfrak{M}$ and $D(TQ) = \{\xi \in \mathcal{H} : Q\xi \in D(T)\}$ is dense in $\mathcal{H}$, then $QTQ \,\eta\, Q\mathfrak{M}Q$.

Let $\mathfrak{M}$ be a von Neumann algebra on a Hilbert space $\mathcal{H}$ and φ a normal faithful semifinite trace defined on $\mathfrak{M}_+$. For each $p \geq 1$, let

$$\mathcal{J}_p = \{X \in \mathfrak{M} : \varphi(|X|^p) < \infty\}.$$

Then $\mathcal{J}_p$ is a *-ideal of $\mathfrak{M}$. Following [17], we denote with $L^p(\varphi)$ the Banach space completion of $\mathcal{J}_p$ with respect to the norm

$$\|X\|_{p,\varphi} := \varphi(|X|^p)^{1/p}, \quad X \in \mathcal{J}_p.$$

One usually defines $L^\infty(\varphi) := \mathfrak{M}$. Thus, if φ is a finite trace, then $L^\infty(\varphi) \subset L^p(\varphi)$ for every $p \geq 1$. As shown in [17], if $X \in L^p(\varphi)$, then X is a measurable operator.

If A is a measurable operator and $A \geq 0$, one defines the *integral* of A by

$$\mu(A) = \sup\{\varphi(X) :\ 0 \leq X \leq A,\ X \in \mathcal{J}_1\}.$$

Then the space $L^p(\varphi)$ can also be defined [17] as the space of all measurable operators A such that $\mu(|A|^p) < \infty$.

The integral of an element $A \in L^p(\varphi)$ can be defined, in obvious way, taking into account that any measurable operator A can be decomposed as $A = B_+ - B_- + iC_+ - iC_-$, where $B = \frac{A+A^*}{2}$ and $C = \frac{A-A^*}{2i}$ and B_+, B_- (resp. C_+, C_-) are the positive and negative parts of B (resp. C).

1.2. **Locally convex quasi C*-algebras.** In what follows we recall some definitions and facts needed in the sequel.

Definition 1.3. Let $\mathfrak{X}$ be a complex vector space and $\mathfrak{A}_0$ a * -algebra contained in $\mathfrak{X}$. $\mathfrak{X}$ is said a *quasi *-algebra with distinguished *-algebra* $\mathfrak{A}_0$ (or, simply, over $\mathfrak{A}_0$) if

(i) the multiplication of $\mathfrak{A}_0$ is extended on $\mathfrak{X}$ as follows: the correspondences
$\mathfrak{X} \times \mathfrak{A}_0 \to \mathfrak{A} : (a, x) \mapsto ax$ (left multiplication of x by a) and
$\mathfrak{A}_0 \times \mathfrak{X} \to \mathfrak{A} : (x, a) \mapsto xa$ (right multiplication of x by a)
are always defined and are bilinear;

(ii) $x_1(x_2a) = (x_1x_2)a, (ax_1)x_2 = a(x_1x_2)$ and $x_1(ax_2) = (x_1a)x_2$, for all $x_1, x_2 \in \mathfrak{A}_0$ and $a \in \mathfrak{X}$;

(iii) the involution $*$ of $\mathfrak{A}_0$ is extended on $\mathfrak{X}$, denoted also by $*$, and satisfies $(ax)^* = x^*a^*$ and $(xa)^* = a^*x^*$, for all $x \in \mathfrak{A}_0$ and $a \in \mathfrak{X}$.

Thus a *quasi *-algebra* [18] is a couple $(\mathfrak{X}, \mathfrak{A}_0)$, where $\mathfrak{X}$ is a vector space with involution *, $\mathfrak{A}_0$ is a *-algebra and a vector subspace of $\mathfrak{X}$ and $\mathfrak{X}$ is an $\mathfrak{A}_0$-bimodule whose module operations and involution extend those of $\mathfrak{A}_0$. The *unit* of $(\mathfrak{X}, \mathfrak{A}_0)$ is an element $e \in \mathfrak{A}_0$ such that $xe = ex = x$, for every $x \in \mathfrak{X}$.

A quasi *-algebra $(\mathfrak{X}, \mathfrak{A}_0)$ is said to be *locally convex* if $\mathfrak{X}$ is endowed with a topology τ which makes $\mathfrak{X}$ a locally convex space and such that the involution $a \mapsto a^*$ and the multiplications $a \mapsto ab$, $a \mapsto ba$, $b \in \mathfrak{A}_0$, are continuous. If τ is a norm topology and the involution is isometric with respect to the norm, we say that $(\mathfrak{X}, \mathfrak{A}_0)$ is a *normed quasi *-algebra* and, if it is complete, we say it is a *Banach quasi*-algebra*.

Let $\mathfrak{A}_0[\|\cdot\|_0]$ be a C*-algebra. We shall use the symbol $\|\cdot\|_0$ of the C*-norm to denote the corresponding topology. Suppose that τ is a topology on $\mathfrak{A}_0$ such that $\mathfrak{A}_0[\tau]$ is a locally convex $*$-algebra. Then, the topologies τ, $\|\cdot\|_0$ on $\mathfrak{A}_0$ are *compatible* whenever each Cauchy net in both topologies that converges with respect to one of them, also converges with respect to the other one.

Under certain conditions on τ a quasi $*$-subalgebra $\mathfrak{A}$ of the quasi $*$-algebra $\mathfrak{X} = \widetilde{\mathfrak{A}_0}[\tau]$ over $\mathfrak{A}_0$ is named *locally convex quasi C*-algebra*. More precisely, let $\{p_\lambda\}_{\lambda\in\Lambda}$ be a directed family of seminorms defining the topology τ. Suppose that τ is compatible with $\|\cdot\|_0$ and that it has the following properties:

- (T_1) $\mathfrak{A}_0[\tau]$ is a locally convex $*$-algebra with separately continuous multiplication.
- (T_2) $\tau \preceq \|\cdot\|_0$.

Then, the identity map $\mathfrak{A}_0[\|\cdot\|_0] \to \mathfrak{A}_0[\tau]$ extends to a continuous $*$-linear map $\mathfrak{A}_0[\|\cdot\|_0] \to \widetilde{\mathfrak{A}_0}[\tau]$. Since $\tau, \|\cdot\|_0$ are compatible, the C*-algebra $\mathfrak{A}_0[\|\cdot\|_0]$ can be regarded as embedded into $\widetilde{\mathfrak{A}_0}[\tau]$. It is easily shown that $\widetilde{\mathfrak{A}_0}[\tau]$ is a quasi $*$-algebra over $\mathfrak{A}_0$ (cf. [13, Section 3]).

We denote by $(\mathfrak{A}_0)_+$ the set of all positive elements of the C*-algebra $\mathfrak{A}_0[\|\cdot\|_0]$.

Further, we employ the following two extra conditions (T_3), (T_4) for the locally convex topology τ on $\mathfrak{A}_0$

- (T_3) For each $\lambda \in \Lambda$, there exists $\lambda' \in \Lambda$ such that
$$p_\lambda(xy) \leq \|x\|_0 p_{\lambda'}(y), \forall\ x, y \in \mathfrak{A}_0 \text{ with } xy = yx;$$
- (T_4) The set $\mathcal{U}(\mathfrak{A}_0)_+ := \{x \in (\mathfrak{A}_0)_+ : \|x\|_0 \leq 1\}$ is τ-closed.

Definition 1.4. By a *locally convex quasi C*-algebra* over $\mathfrak{A}_0$, (see [3]), we mean any quasi $*$-subalgebra $\mathfrak{A}$ of the locally convex quasi $*$-algebra $\mathfrak{X} = \widetilde{\mathfrak{A}_0}[\tau]$ over $\mathfrak{A}_0$, where $\mathfrak{A}_0[\|\cdot\|_0]$ is a C*-algebra with identity e and τ a locally convex topology on $\mathfrak{A}_0$, defined by a directed family of seminorms $\{p_\lambda\}_{\lambda\in\Lambda}$, satisfying the conditions (T_1)-(T_4).

The following examples have been discussed in [3].

Example 1.5. [CQ*-algebras] Let $\mathfrak{A}_0$ be a C^*-algebra, with norm $\|\cdot\|$ and involution *. Let $\|\cdot\|_1$ be a norm on $\mathfrak{A}_0$, weaker than $\|\cdot\|$ and such that, for every $a, b \in \mathfrak{A}$

(i) $\|ab\|_1 \leq \|a\|_1\|b\|$,
(ii) $\|a^*\|_1 = \|a\|_1$.

Let $\mathfrak{X}$ denote the $\|\cdot\|_1$-completion of $\mathfrak{A}_0$; then [1] the couple $(\mathfrak{X}, \mathfrak{A}_0)$ is called a *CQ*-algebra*. Every CQ*-algebra is a locally convex quasi C*-algebra.

Example 1.6. The space $L^p([0,1])$, with $1 \leq p < +\infty$ is a Banach $L^\infty([0,1])$-bimodule. The couple $(L^p([0,1]), L^\infty([0,1])$ may be regarded as a CQ*-algebra thus a locally convex quasi C*-algebra over $L^\infty([0,1])$.

2. Locally convex quasi C*-algebras of measurable operators

Let $\mathfrak{M}$ be a von Neumann algebra on a Hilbert space $\mathcal{H}$ and φ a normal faithful semifinite trace on $\mathfrak{M}_+$, then, as shown in [9], $(L^p(\varphi), L^\infty(\varphi) \cap L^p(\varphi))$ is a *Banach quasi*-algebra* and if φ is a finite trace, $(L^p(\varphi), \mathfrak{M})$ is a CQ*-algebra.

In analogy to [9] we consider the following two sets of sesquilinear forms enjoying certain invariance properties.

Definition 2.1. Let $(\mathfrak{X}, \mathfrak{A}_0)$ be a locally convex quasi C*-algebra with unit e. We denote by $\mathcal{S}(\mathfrak{X})$ the set of all sesquilinear forms Ω on $\mathfrak{X} \times \mathfrak{X}$ with the following properties:

(i) $\Omega(x,x) \geq 0 \quad \forall x \in \mathfrak{X}$;
(ii) $\Omega(xa, b) = \Omega(a, x^*b) \quad \forall x \in \mathfrak{X}, \ \forall a, b \in \mathfrak{A}_0$;
(iii) $|\Omega(x,y)| \leq p(x)p(y)$ for some τ-continuous seminorm p on $\mathfrak{X}$ and all $x, y \in \mathfrak{X}$;
(iv) $\Omega(e,e) \leq 1$.

The locally convex quasi C*-algebra $(\mathfrak{X}, \mathfrak{A}_0)$ is called **-semisimple* if $x \in \mathfrak{X}$, $\Omega(x,x) = 0$, for every $\Omega \in \mathcal{S}(\mathfrak{X})$, implies $x = 0$.

We denote by $\mathcal{T}(\mathfrak{X}) \subseteq \mathcal{S}(\mathfrak{X})$ the set of all sesquilinear forms $\Omega \in \mathcal{S}(\mathfrak{X})$ with the following property

(iv) $\Omega(x,x) = \Omega(x^*, x^*), \ \forall x \in \mathfrak{X}$.

Remark 2.2. Notice that

- By (iv) of Definition 2.1 and by polarization, we get
$$\Omega(y^*, x^*) = \Omega(x,y), \quad \forall x, y \in \mathfrak{X}.$$
- The set $\mathcal{T}(\mathfrak{X})$ is convex.

Example 2.3. Let $\mathfrak{M}$ be a von Neumann algebra and φ a normal faithful semifinite trace on $\mathfrak{M}_+$. Then, $(L^p(\varphi), \mathcal{J}_p)$, $p \geq 2$, is a *-semisimple Banach quasi *-algebra. If φ is a finite trace (we assume $\varphi(\mathbb{I}) = 1$), then $(L^p(\varphi), \mathfrak{M})$, with $p \geq 2$, is a *-semisimple locally convex quasi C*-algebra. If $p \geq 2$, L^p-spaces possess a sufficient family of positive sesquilinear forms. Indeed, in this case, since for every $W \in L^p(\varphi)$, $|W|^{p-2} \in L^{p/(p-2)}(\varphi)$, then the sesquilinear form Ω_W defined by

$$\Omega_W(X,Y) := \frac{\varphi[X(Y|W|^{p-2})^*]}{\| W \|_{p,\varphi}^{p-2}}$$

is positive and satisfies the conditions (i),(ii),(iii) and (iv) of Definition 2.1, (see [9], and [24] for more details). Moreover,

$$\Omega_W(W,W) = \| W \|_{p,\varphi}^p .$$

[1] In previous papers this structure was called *proper* CQ*-algebra. Since this is the sole case we consider here we will systematically omit the specification *proper*.

Remark 2.4. The notion of *-semisimplicity of locally convex partial *-algebras has been studied in full generality in [2] and [14].

Definition 2.5. Let $\mathfrak{M}$ be a von Neumann algebra and φ a normal faithful semifinite trace defined on $\mathfrak{M}_+$. We denote by $L^p_{\rm loc}(\varphi)$ the set of all measurable operators T such that $TP \in L^p(\varphi)$, for every central φ-finite projection P of $\mathfrak{M}$.

Remark 2.6. The von Neumann algebra $\mathfrak{M}$ is a subset of $L^p_{\rm loc}(\varphi)$. Indeed, if $X \in \mathfrak{M}$, then for every φ-finite central projection P of $\mathfrak{M}$ the product XP belongs to the *-ideal $\mathcal{J}_p$.

Throughout this section we are given a von Neumann algebra $\mathfrak{M}$ on a Hilbert space $\mathcal{H}$ with a family $\{P_j\}_{j\in J}$ of φ-finite central projections of $\mathfrak{M}$ such that

- if $l, m \in J$, $l \neq m$, then $P_l P_m = 0$ (i.e., the P_j's are orthogonal);
- $\bigvee_{j\in J} P_j = \mathbb{I}$; where $\bigvee_{j\in J} P_j$ denotes the projection onto the subspace generated by $\{P_j\mathcal{H} : j \in J\}$.

The previous two conditions are always realized in a von Neumann $\mathfrak{M}$ algebra with a faithful normal semifinite trace (see Lemma 1.1 and [15, 20] for more details).

If φ is a normal faithful semifinite trace on $\mathfrak{M}_+$, we define, for each $X \in \mathfrak{M}$, the following seminorms $q_j(X) := \|XP_j\|_{p,\varphi}$ on $\mathfrak{M}$, $j \in J$. The traslation-invariant locally convex topology defined by the system $\{q_j : j \in J\}$ is denoted by τ_p.

Definition 2.7. Let $\mathfrak{M}$ be a von Neumann algebra and φ a normal faithful semifinite trace defined on $\mathfrak{M}_+$. We denote by $\widetilde{\mathfrak{M}}^{\tau_p}$ the τ_p completion of $\mathfrak{M}$.

Proposition 2.8. *Let $\mathfrak{M}$ be a von Neumann algebra and φ a normal faithful semifinite trace on $\mathfrak{M}_+$. Then $L^p_{\rm loc}(\varphi) \subseteq \widetilde{\mathfrak{M}}^{\tau_p}$. Moreover, if there exists a family $\{P_j\}_{j\in J}$ as above, where all P_j' s are mutually equivalent, then $L^p_{\rm loc}(\varphi) = \widetilde{\mathfrak{M}}^{\tau_p}$.*

Proof. From Remark 2.6, $\mathfrak{M} \subseteq L^p_{\rm loc}(\varphi)$. If $Y \in L^p_{\rm loc}(\varphi)$, for every $j \in J$ we have $YP_j \in L^p(\varphi)$. Then, for every $j \in J$, there exist $(X^j_n)_{n=1}^\infty \subseteq \mathcal{J}_p$ such that $\|X^j_n - YP_j\|_{p,\varphi} \underset{n\to\infty}{\longrightarrow} 0$.

Let $\mathbb{F}_J$ be the family of finite subsets of J ordered by inclusion and $F \in \mathbb{F}_J$.

We put

$$T_{n,F} := \sum_{j\in F} X^j_n P_j \in \mathfrak{M}.$$

Then the net $(T_{n,F})$ converges to Y with respect to τ_p. Indeed, for every $m \in J$,

$$q_m(T_{n,F} - Y) = \|(T_{n,F} - Y)P_m\|_{p,\varphi} = \|(X^m_n - Y)P_m\|_{p,\varphi}$$

for sufficiently large F. Thus, the inequality $\|(X^m_n - Y)P_m\|_{p,\varphi} \leq \|X^m_n - YP_m\|_{p,\varphi}$ implies that $q_m(T_{n,F} - Y) \underset{n,F}{\longrightarrow} 0$.

Hence $L^p_{\rm loc}(\varphi) \subseteq \widetilde{\mathfrak{M}}^{\tau_p}$.

On the other hand, assume that all P_j' s are mutually equivalent. Then, if $Y \in \widetilde{\mathfrak{M}}^{\tau_p}$, there exists a net $(X_\alpha) \subseteq \mathfrak{M}$ such that $X_\alpha \to Y$ with respect to τ_p; hence

$$X_\alpha P_j \to YP_j \in L^p(\varphi) \quad \text{in } \|\cdot\|_{p,\varphi}\,. \tag{1}$$

But for each central $\varphi-$finite projection P we have

$$\varphi(P) = \varphi(P\sum_{j\in J} P_j) = \sum_{j\in J} \varphi(PP_j). \tag{2}$$

By our assumption, for any $l, m \in J$, we may pick $U \in \mathfrak{M}$ so that $U^*U = P_l$ and $UU^* = P_m$, hence,

$$\varphi(PP_l) = \varphi(PU^*U) = \varphi(UPU^*) = \varphi(PUU^*) = \varphi(PP_m).$$

So, all terms on the right hand side of (2) are equal and since the above series converges, only a finite number of them can be nonzero. Thus, for some $s \in \mathbb{N}$ we may write $J = \{1, \ldots, s\}$ and then

$$P = P\sum_{j\in J} P_j = P\sum_{j=1}^{s} P_j = \sum_{j=1}^{s} PP_j \tag{3}$$

and hence

$$YP = \sum_{j=1}^{s} YPP_j = \sum_{j=1}^{s} YP_jP \in L^p(\varphi). \tag{4}$$

Therefore, if $Y \in \widetilde{\mathfrak{M}}^{\tau_p}$, for each central $\varphi-$finite projection P, we have $YP \in L^p(\varphi)$. Hence $L^p_{\mathrm{loc}}(\varphi) \supseteq \widetilde{\mathfrak{M}}^{\tau_p}$.

□

Remark 2.9. In general it is not guaranteed that a von Neumann algebra possesses an orthogonal family $\{P_j\}_{j\in J}$ of mutually equivalent finite central projection such that $\bigvee_{j\in J} P_j = \mathbb{I}$, but, if this is the case, then $L^p_{\mathrm{loc}}(\varphi) = \widetilde{\mathfrak{M}}^{\tau_p}$.

Theorem 2.10. *Let $\mathfrak{M}$ be a von Neumann algebra on a Hilbert space $\mathcal{H}$ and φ a normal faithful semifinite trace on $\mathfrak{M}_+$. Then $(\widetilde{\mathfrak{M}}^{\tau_p}, \mathfrak{M})$ is a locally convex quasi C*-algebra with respect to τ_p, consisting of measurable operators.*

Proof. The topology τ_p satisfies the properties (T_1)-(T_4). We just prove here (T_3)-(T_4).

- (T_3) For each $\lambda \in J$,

$$q_\lambda(XY) = \| P_\lambda XY \|_{p,\varphi} \leq \|X\| \, \| P_\lambda Y \|_{p,\varphi} = \|X\| q_\lambda(Y), \quad \forall X, Y \in \mathfrak{M};$$

- (T_4) The set $\mathcal{U}(\mathfrak{M})_+ := \{X \in (\mathfrak{M})_+ : \|X\| \leq 1\}$ is τ_p-closed. To see this consider a net $\{F_\alpha\}$ in $\mathcal{U}(\mathfrak{M})_+ : F_\alpha \to F$ in the topology τ_p, then for each $j \in J$, we have $\| (F_\alpha - F)P_j \|_{p,\varphi} \to 0$. By assumption on P_j, the trace φ is a normal faithful finite trace on the von Neumann algebra on $P_j\mathfrak{M}_+$. Then (see [9]) $(L^p(\varphi), P_j\mathfrak{M})$ is a CQ*-algebra. Therefore, using (T_4) for $(L^p(\varphi), P_j\mathfrak{M})$, we have $FP_j \in \mathcal{U}(P_j\mathfrak{M})_+$ for each $j \in J$. This, by definition, implies that $F \in \mathfrak{M}$. Indeed, for every

$$h = \sum_{j\in J} P_j h \in \mathcal{H} = \bigoplus_{j\in J} P_j\mathcal{H}$$

we have

$$\begin{aligned} \| Fh \|^2_{\mathcal{H}} &= \sum_{j\in J} \| FP_jh \|^2 \\ &= \sum_{j\in J} \| FP_jP_jh \|^2 \leq \sum_{j\in J} \| P_jh \|^2 \\ &= \| h \|^2_{\mathcal{H}} . \end{aligned}$$

Hence $F \in \mathcal{U}(\mathfrak{M})_+$.

□

Remark 2.11. By Proposition 2.8, $(L^p_{\rm loc}(\varphi), \mathfrak{M})$ itself is a *locally convex quasi C*-algebra* with respect to τ_p.

3. The representation theorems

Let $(\mathfrak{X}, \mathfrak{A}_0)$ be a locally convex quasi C*-algebra with a unit e. For each $\Omega \in \mathcal{T}(\mathfrak{X})$, we define a linear functional ω_Ω on $\mathfrak{A}_0$ by

$$\omega_\Omega(a) := \Omega(a, e) \qquad a \in \mathfrak{A}_0.$$

We have

$$\omega_\Omega(a^*a) = \Omega(a^*a, e) = \Omega(a, a) = \Omega(a^*, a^*) = \omega_\Omega(aa^*) \geq 0.$$

This shows at once that ω_Ω is positive and tracial.

By the Gelfand-Naimark theorem each C^*-algebra is isometrically *-isomorphic to a C^*-algebra of bounded operators in Hilbert space. This isometric *-isomorphism is called the *universal *-representation*. We denote it by π.

For every $\Omega \in \mathcal{T}(\mathfrak{X})$ and $a \in \mathfrak{A}_0$, we put

$$\varphi_\Omega(\pi(a)) = \omega_\Omega(a).$$

Then, for each $\Omega \in \mathcal{T}(\mathfrak{X})$, φ_Ω is a positive bounded linear functional on the operator algebra $\pi(\mathfrak{A}_0)$.

Clearly,

$$\varphi_\Omega(\pi(a)) = \omega_\Omega(a) = \Omega(a, e).$$

Using the fact that the family $\{p_\lambda\}$ is directed, we conclude that there exist $\gamma > 0$ and $\lambda \in \Lambda$ such that

$$| \varphi_\Omega(\pi(a)) | = | \omega_\Omega(a) | = | \Omega(a, e) | \leq \gamma^2 p_\lambda(ae) p_\lambda(e).$$

Then, by (T_3), for some $\lambda' \in \Lambda$

$$| \varphi_\Omega(\pi(a)) | \leq \gamma^2 \|a\|_0 p_{\lambda'}(e)^2.$$

Thus φ_Ω is continuous on $\pi(\mathfrak{A}_0)$.

By [15, Vol. 2, Proposition 10.1.1], φ_Ω is weakly continuous and so it extends uniquely to $\pi(\mathfrak{A}_0)''$, by the Hahn-Banach theorem. Moreover, since φ_Ω is a trace on $\pi(\mathfrak{A}_0)$, the extension $\widetilde{\varphi_\Omega}$ is also a trace on the von Neumann algebra $\mathfrak{M} := \pi(\mathfrak{A}_0)''$ generated by $\pi(\mathfrak{A}_0)$.

Clearly, the set $\mathfrak{N}_\mathcal{T}(\mathfrak{A}_0) = \{\widetilde{\varphi_\Omega};\, \Omega \in \mathcal{T}(\mathfrak{X})\}$ is convex.

Definition 3.1. The locally convex quasi C*-algebra $(\mathfrak{X}, \mathfrak{A}_0)$ is said to be *strongly *-semisimple* if

(a) the equality $\Omega(x,x) = 0$, for every $\Omega \in \mathcal{T}(\mathfrak{X})$, implies $x = 0$;
(b) the set $\mathfrak{N}_{\mathcal{T}}(\mathfrak{A}_0)$ is w^*-closed.

Remark 3.2. If $(\mathfrak{X}, \mathfrak{A}_0)$ is a *CQ*-algebra*, by [9, Proposition 4.1], (b) is automatically satisfied.

Example 3.3. Let $\mathfrak{M}$ be a von Neumann algebra and φ a normal faithful semifinite trace on $\mathfrak{M}_+$. Then, as seen in Example 2.3, if φ is a finite trace, then $(L^p(\varphi), \mathfrak{M})$, with $p \geq 2$, is a *-semisimple locally convex quasi C*-algebra. The conditions (a) and (b) of Definition 3.1 are satisfied. Indeed, in this case, the set $\mathfrak{N}_{\mathcal{T}}(\mathfrak{M})$ is w^*-closed by [9, Proposition 4.1]. Therefore $(L^p(\varphi), \mathfrak{M})$, with φ finite, is a strongly *-semisimple locally convex quasi C*-algebra .

Let $(\mathfrak{X}, \mathfrak{A}_0)$ be a locally convex quasi C*-algebra with unit e, π the universal representation of $\mathfrak{A}_0$ and $\mathfrak{M} = \pi(\mathfrak{A}_0)''$. Denote by $\| f \|^\sharp$ the norm of a bounded functional f on $\mathfrak{M}$ and by $\mathfrak{M}^\sharp$ the topological dual of $\mathfrak{M}$ then, the norm $\| \widetilde{\varphi_\Omega} \|^\sharp$ of $\widetilde{\varphi_\Omega}$ as a linear functional on $\mathfrak{M}$ equals the norm of φ_Ω as a functional on $\pi(\mathfrak{A}_0)$.

By (iv) of Definition 2.1, $\| \widetilde{\varphi_\Omega} \|^\sharp = \widetilde{\varphi}_\Omega(\pi(e)) = \Omega(e,e) \leq 1$.

Hence, if (b) of Definition 3.1 is satisfied, the set $\mathfrak{N}_{\mathcal{T}}(\mathfrak{A}_0)$, being a w^*-closed subset of the unit ball of $\mathfrak{M}^\sharp$, is w^*-compact.

Let $\mathfrak{EN}_{\mathcal{T}}(\mathfrak{A}_0)$ be the set of extreme points of $\mathfrak{N}_{\mathcal{T}}(\mathfrak{A}_0)$; then $\mathfrak{N}_{\mathcal{T}}(\mathfrak{A}_0)$ coincides with w^*-closure of the convex hull of $\mathfrak{EN}_{\mathcal{T}}(\mathfrak{A}_0)$. Let $\pi : \mathfrak{A}_0 \hookrightarrow \mathcal{B}(\mathcal{H})$ be its universal representation.

Then $\mathfrak{EN}_{\mathcal{T}}(\mathfrak{A}_0)$ is a family of normal finite traces on the von Neumann algebra $\mathfrak{M}$.

We put $\mathcal{F} := \{\Omega \in \mathcal{T}(\mathfrak{X});\ \widetilde{\varphi_\Omega} \in \mathfrak{EN}_{\mathcal{T}}(\mathfrak{A}_0)\}$ and denote by P_Ω the support projection corresponding to the trace $\widetilde{\varphi_\Omega}$. By [9, Lemma 3.5], $\{P_\Omega\}_{\Omega\in\mathcal{F}}$ consists of mutually orthogonal projections and if $Q := \bigvee_{\Omega\in\mathcal{F}} P_\Omega$ then

$$\mu = \sum_{\widetilde{\varphi_\Omega}\in\mathfrak{EN}_{\mathcal{T}}(\mathfrak{A}_0)} \widetilde{\varphi_\Omega}$$

is a normal faithful semifinite trace defined on the direct sum (see [20] and [26]) of von Neumann algebras

$$Q\mathfrak{M} = \bigoplus_{\Omega\in\mathcal{F}} P_\Omega \mathfrak{M}.$$

Theorem 3.4. *Let $(\mathfrak{X}, \mathfrak{A}_0)$ be a strongly *-semisimple locally convex quasi C*-algebra with unit e, π the universal representation of $\mathfrak{A}_0$. Then there exists a monomorphism*

$$\Phi : x \in \mathfrak{X} \to \Phi(x) := \widetilde{X} \in \widetilde{Q\mathfrak{M}}^{\tau_2}$$

with the following properties:

(i) Φ extends the isometry $\pi : \mathfrak{A}_0 \hookrightarrow \mathcal{B}(\mathcal{H})$ given by the Gelfand-Naimark theorem;
(ii) $\Phi(x^*) = \Phi(x)^*$, for every $x \in \mathfrak{X}$;
(iii) $\Phi(xy) = \Phi(x)\Phi(y)$, for all $x, y \in \mathfrak{X}$ such that $x \in \mathfrak{A}_0$ or $y \in \mathfrak{A}_0$.

Proof. Let $\{p_\lambda\}_{\lambda\in\Lambda}$ be, as before, the family of seminorms defining the topology τ of $\mathfrak{X}$. For fixed $x \in \mathfrak{X}$, there exists a net $\{a_\alpha;\, \alpha \in \Delta\}$ of elements of $\mathfrak{A}_0$ such that $p_\lambda(a_\alpha - x) \to 0$ for each $\lambda \in \Lambda$. We put $X_\alpha = \pi(a_\alpha)$.

By (iii) of Definition 2.1, for every $\Omega \in \mathcal{T}(\mathfrak{X})$, there exist $\gamma > 0$ and $\lambda' \in \Lambda$ such that for each $\alpha, \beta \in \Delta$

$$\begin{aligned}\|P_\Omega(X_\alpha - X_\beta)\|_{2,\widetilde{\varphi_\Omega}} &= \|P_\Omega(\pi(a_\alpha) - \pi(a_\beta))\|_{2,\widetilde{\varphi_\Omega}} \\ &= [\widetilde{\varphi_\Omega}(|P_\Omega(\pi(a_\alpha) - \pi(a_\beta))|^2)]^{1/2} = \\ &= [\Omega\left((a_\alpha - a_\beta)^*(a_\alpha - a_\beta), e\right)]^{1/2} \\ &= [\Omega(a_\alpha - a_\beta, a_\alpha - a_\beta)]^{1/2} \leq \gamma p_{\lambda'}(a_\alpha - a_\beta) \underset{\alpha,\beta}{\longrightarrow} 0.\end{aligned}$$

Let $\widetilde{X_\Omega}$ be the $\|\cdot\|_{2,\widetilde{\varphi_\Omega}}$-limit of the net $(P_\Omega X_\alpha)$ in $L^2(\widetilde{\varphi_\Omega})$. Clearly $\widetilde{X_\Omega} = P_\Omega \widetilde{X_\Omega}$

We define

$$\Phi(x) := \sum_{\Omega\in\mathcal{F}} P_\Omega \widetilde{X_\Omega} =: \widetilde{X}.$$

Clearly $\widetilde{X} \in \widetilde{Q\mathfrak{M}}^{\tau_2}$.

It is easy to see that the map $x \ni \mathfrak{X} \mapsto \widetilde{X} \in \widetilde{Q\mathfrak{M}}^{\tau_2}$ is well defined and injective. Indeed, if $a_\alpha \to 0$, there exist $\gamma > 0$ and $\lambda' \in \Lambda$ such that

$$\begin{aligned}\|P_\Omega X_\alpha\|_{2,\widetilde{\varphi_\Omega}} &= \|P_\Omega \pi(a_\alpha)\|_{2,\widetilde{\varphi_\Omega}} \\ &= [\widetilde{\varphi_\Omega}(|P_\Omega(\pi(a_\alpha)|^2)]^{1/2} = \\ &= [\Omega\left(a_\alpha^* a_\alpha, e\right)]^{1/2} \\ &= [\Omega(a_\alpha, a_\alpha)]^{1/2} \leq \gamma p_{\lambda'}(a_\alpha) \to 0.\end{aligned}$$

Thus $P_\Omega(X_\alpha) = 0$ for every $\Omega \in \mathcal{T}(\mathfrak{X})$, then $\widetilde{X} = 0$. Moreover if $P_\Omega \widetilde{X} = 0$, for each $\Omega \in \mathcal{F}$, then $\Omega(x, x) = 0$ for every $\Omega \in \mathcal{F}$. Since every $\Omega \in \mathcal{T}(\mathfrak{X})$ is a w*-limit of of convex combinations of elements of $\mathcal{F}$, we get $\Omega(x, x) = 0$ for every $\Omega \in \mathcal{T}(\mathfrak{X})$. Therefore, by assumption, $x = 0$.

□

Remark 3.5.

In the same way we can prove that

- If $(\mathfrak{X}, \mathfrak{A}_0)$ is a strongly *-semisimple locally convex quasi C*-algebra and there exists a faithful $\Omega \in \mathcal{T}(\mathfrak{X})$ (i.e., the equality $\Omega(x, x) = 0,$ implies $x = 0$) then there exists a monomorphism

  $$\Phi : x \in \mathfrak{X} \to \Phi(x) := \widetilde{X} \in L^2(\widetilde{\varphi_\Omega})$$

  with the following properties:

  (i) Φ extends the isometry $\pi : \mathfrak{A}_0 \hookrightarrow \mathcal{B}(\mathcal{H})$ given by the Gelfand-Naimark theorem;

  (ii) $\Phi(x^*) = \Phi(x)^*,$ for every $x \in \mathfrak{X}$

  (iii) $\Phi(xy) = \Phi(x)\Phi(y)$, for all $x, y \in \mathfrak{X}$ such that $x \in \mathfrak{A}_0$ or $y \in \mathfrak{A}_0$.

- If the semifinite von Neumann algebra $\pi(\mathfrak{A}_0)''$ admits an orthogonal family of mutually equivalent projections $\{P_i'; i \in I\}$ such that $\sum_{i\in I} P_i' = \mathbb{I}$, then is easy to see that the map $x \in \mathfrak{X} \to \widetilde{X} \in L^2_{loc}(\tau)$ is a monomorphism.

Acknowledgement. The authors wish to express their gratitude to the referee for pointing out several inaccuracies in a previous version of the paper and for his/her fruitful comments.

References

1. J.-P. Antoine, A.Inoue and C.Trapani, *Partial *-algebras and their Operator Realizations*, Kluwer, Dordrecht, 2002.
2. J-P. Antoine, G. Bellomonte, C. Trapani, *Fully representable and *-semisimple topological partial *-algebras*, Studia Mathematica, No.2, **208** (2012), 167-194 DOI: 10.4064/sm208-2-4.
3. F. Bagarello, M. Fragoulopoulou, A. Inoue and C. Trapani, *Locally convex quasi C*-normed algebras. J. Math. Anal. Appl.*, **366**, (2010), no. 2, 593606.
4. F.Bagarello, A.Inoue and C.Trapani, *Some classes of topological quasi *-algebras*, Proc. Amer. Math. Soc., **129** (2001), 2973-2980.
5. F.Bagarello and C.Trapani,*States and representations of CQ* -algebras*, Ann. Inst. H. Poincaré **61**, 103-133(1994).
6. F.Bagarello and C.Trapani, *L^p-spaces as quasi *-algebras*, J.Math. Anal. Appl. **197**, 810-824 (1996).
7. F.Bagarello and C.Trapani, *CQ*-algebras: structure properties* Publ. RIMS, Kyoto Univ. **32**, 85-116 (1996).
8. F.Bagarello and C.Trapani, *Morphisms of certain Banach C*-modules*, Publ. RIMS, Kyoto Univ. **36** (2000) 681-705
9. F. Bagarello, C. Trapani and S. Triolo, *Quasi $*$-algebras of measurable operators*, Studia Mathematica **172** (2006), no. 3, 289–305.
10. F. Bagarello, C. Trapani and S. Triolo, *A note on faithful traces on a von Neumann algebra*, Rend. Circ. Mat. Palermo **55** (2006), no. 1, 21–28.
11. F. Bagarello, C. Trapani, S. Triolo; *Representable states on quasi-local quasi *-algebras*, J. Math. Phys. 52 (2011), no. 1, 013510, 11 pp.
12. B. Bongiorno, C. Trapani, S. Triolo *Extensions of positive linear functionals on a topological *-algebra*, Rocky Mt. J. Math. 40(6), 17451777 (2010).
13. M. Fragoulopoulou, A. Inoue and K.-D. Kürsten, *On the completion of a C*-normed algebra under a locally convex algebra topology*, Contemporary Math. Vol.5, **427**; 155-166 (2007).
14. M. Fragoulopoulou, C. Trapani, C. and S. Triolo, *Locally convex quasi *-algebras with sufficiently many *-representations*, J.Math. Anal. Appl., **388** (2012), 1180-1193 .
15. R. V. Kadison and J. R. Ringrose, *Fundamentals of the Theory of Operator Algebras*, Vol. 1-4, Academic Press, Orlando, London, 1986.
16. C. La Russa and S. Triolo; *Radon - Nikodym theorem in topological quasi *-algebras*. J. Operator Th. **69**, (2013) 423–433.
17. E. Nelson, *Notes on noncommutative integration*, J. Funct. Anal., **15** (1974) 103-116
18. K. Schmüdgen, *Unbounded Operator Algebras and Representation Theory*, Akademie-Verlag, Berlin, 1990.
19. I. E. Segal, *A noncommutative extension of abstract integration*, *Ann. Math.* **57** (1953), 401–457.
20. M. Takesaki, *Theory of Operator Algebras.I*, Springer-Verlag, New York, 1979.
21. C.Trapani, *CQ*-algebras of operators and application to Quantum Models*, in Proceedings of the Second ISAAC Conference, Fukuoka, Japan, Volume 1, 679-685, Kluwer (2000).
22. C. Trapani and S. Triolo, *Representations of modules over a $*$-algebra and related seminorms*, Studia Mathematica **184** (2008), 133–148.
23. C. Trapani and S. Triolo, *Faithful Representations of left C*-modules*, Rend. Circolo Mat. Palermo 59 (2010), 295-302.

24. S. Triolo, *WQ*-Algebras of measurable operators*, Indian J. Pure Appl. Math. **43** (2012), 601-617.
25. S. Triolo, *Moduli di Banach su C* algebre, Rappresentazioni Hilbertiane ed in spazi Lp non commutativi*, Bollettino Umi sez. A Fascicolo **N. 2** - (Agosto 2006).
26. S. Triolo, *Possible extensions of the noncommutative integral*, Rend. Circ. Mat. Palermo, (2) 60 (2011), no 3, 409–416. DOI 10.1007/s12215-011-0063-1.
27. S. Triolo, *Extensions of the non commutative integration*, submitted to Complex Analysis and Operator Theory.

Dipartimento di Matematica e Informatica, Università di Palermo, I-90123 Palermo (Italy)
E-mail address: camillo.trapani@unipa.it

Dipartimento DEIM, Università di Palermo, I-90123 Palermo (Italy)
E-mail address: salvatore.triolo@unipa.it